

# A Remark on the Kolmogorov–Stein Inequality\*

Ha Huy Bang<sup>†</sup>

*Institute of Mathematics, P.O. Box 631, Bo Ho, 10000 Hanoi, Vietnam*

*Submitted by J. L. Brenner*

etaddata, citation and similar papers at [core.ac.uk](https://core.ac.uk)

In this paper, essentially developing the Stein method, we prove the Kolmogorov–Stein inequality for any Orlicz norm (with the same constants). © 1996 Academic Press, Inc.

## 1. INTRODUCTION

A. N. Kolmogorov has given the following result [1]: Let  $f(x)$ ,  $f'(x), \dots, f^{(n)}(x)$  be continuous and bounded on  $\mathbb{R}$ . Then

$$\|f^{(k)}\|_{\infty}^n \leq C_{k,n} \|f\|_{\infty}^{n-k} \|f^{(n)}\|_{\infty}^k,$$

where  $0 < k < n$ ,  $C_{k,n} = K_{n-k}^n / K_n^{(n-k)}$ ,

$$K_i = \frac{4}{\pi} \sum_{j=0}^{\infty} (-1)^j / (2j+1)^{i+1}$$

for even  $i$ , while

$$K_i = \frac{4}{\pi} \sum_{j=0}^{\infty} 1 / (2j+1)^{i+1}$$

for odd  $i$ . Moreover the constants are best possible.

This result has been extended by E. M. Stein to the  $L_p$ -norm [2]. The Kolmogorov–Stein inequality and its variants are a problem of interest for

\*Supported by the National Basic Research Program in Natural Science and by the NCNST “Applied Mathematics.”

<sup>†</sup>E-mail address: hhhbang@thevinh.ac.vn.

many mathematicians and have various applications (see, for example, [3, 4] and their references).

In this paper, essentially developing the Stein method [2], we prove this inequality for an arbitrary Orlicz norm  $\|\cdot\|_\Phi$ . The obtained result has been successfully applied to proving the corresponding imbedding theorems [5–7] for Sobolev–Orlicz spaces of infinite order and the result [8] for any Orlicz norm.

## 2. RESULTS

Let  $\Phi(t): [0, +\infty) \rightarrow [0, +\infty]$  be an arbitrary Young function [9–12], i.e.,  $\Phi(0) = 0$ ,  $\Phi(t) \geq 0$ ,  $\Phi(t) \equiv 0$ , and  $\Phi(t)$  is convex. Denote by

$$\bar{\Phi}(t) = \sup_{s \geq 0} \{ts - \Phi(s)\},$$

which is the Young function conjugate to  $\Phi(t)$  and  $L_\Phi(\mathbb{R})$ , the space of measurable functions  $u(x)$  such that

$$|\langle u, v \rangle| = \left| \int u(x)v(x)dx \right| < \infty$$

for all  $v(x)$  with  $\rho(v, \bar{\Phi}) < \infty$ , where

$$\rho(v, \bar{\Phi}) = \int \bar{\Phi}(|v(x)|) dx.$$

Then  $L_\Phi(\mathbb{R})$  is a Banach space with respect to the Orlicz norm

$$\|u\|_\Phi = \sup_{\rho(v, \bar{\Phi}) \leq 1} \left| \int u(x)v(x) dx \right|,$$

which is equivalent to the Luxemburg norm

$$\|f\|_{(\Phi)} = \inf \left\{ \lambda > 0: \int \Phi(|f(x)|/\lambda) dx \leq 1 \right\} < \infty.$$

We have the following results [9]:

LEMMA 1. *Let  $u \in L_\Phi(\mathbb{R})$  and  $v \in L_{\bar{\Phi}}(\mathbb{R})$ . Then*

$$\int |u(x)v(x)| dx \leq \|u\|_\Phi \|v\|_{(\bar{\Phi})}.$$

LEMMA 2. *Let  $u \in L_\Phi(\mathbb{R})$  and  $v \in L_1(\mathbb{R})$ . Then*

$$\|u * v\|_\Phi \leq \|u\|_\Phi \|v\|_1.$$

Recall that  $\|\cdot\|_\Phi = \|\cdot\|_p$  when  $1 \leq p < \infty$  and  $\Phi(t) = t^p$ ; and  $\|\cdot\|_\Phi = \|\cdot\|_\infty$  when  $\Phi(t) = 0$  for  $0 \leq t \leq 1$  and  $\Phi(t) = \infty$  for  $t > 1$ .

**THEOREM 1.** *Let  $\Phi(t)$  be an arbitrary Young function,  $f(x)$  and its generalized derivative  $f^{(n)}(x)$  be in  $L_\Phi(\mathbb{R})$ . Then  $f^{(k)}(x) \in L_\Phi(\mathbb{R})$  for all  $0 < k < n$  and*

$$\|f^{(k)}\|_\Phi^n \leq C_{k,n} \|f\|_\Phi^{n-k} \|f^{(n)}\|_\Phi^k. \quad (1)$$

*Proof.* We begin to prove (1) with the assumption that  $f^{(k)}(x) \in L_\Phi(\mathbb{R})$ ,  $0 \leq k \leq n$ .

Fix  $0 < k < n$ . It is known that  $\rho(v, \overline{\Phi}) = 1$  if and only if  $\|v\|_{(\overline{\Phi})} = 1$ . Therefore, by the definition we get

$$\|f^{(k)}\|_\Phi = \sup_{\|v\|_{(\overline{\Phi})} \leq 1} \left| \int_{-\infty}^{\infty} f^{(k)}(x) v(x) dx \right|.$$

Let  $\epsilon > 0$ . We choose a function  $v_\epsilon(x) \in L_{\overline{\Phi}}(\mathbb{R})$  such that  $\|v_\epsilon\|_{(\overline{\Phi})} = 1$  and

$$\left| \int_{-\infty}^{\infty} f^{(k)}(x) v_\epsilon(x) dx \right| \geq \|f^{(k)}\|_\Phi - \epsilon. \quad (2)$$

Put

$$F_\epsilon(x) = \int_{-\infty}^{\infty} f(x+y) v_\epsilon(y) dy.$$

Then  $F_\epsilon(x) \in L_\infty(\mathbb{R})$  by virtue of Lemma 1, and

$$F_\epsilon^{(r)}(x) = \int_{-\infty}^{\infty} f^{(r)}(x+y) v_\epsilon(y) dy, \quad 0 \leq r \leq n. \quad (3)$$

Actually, for every function  $\varphi(x) \in C_0^\infty(\mathbb{R})$  it follows from the assumption and Lemma 1 that

$$\begin{aligned} \langle F_\epsilon^{(r)}(x), \varphi(x) \rangle &= (-1)^r \langle F_\epsilon(x), \varphi^{(r)}(x) \rangle \\ &= (-1)^r \int_{-\infty}^{\infty} \left( \int_{-\infty}^{\infty} f(x+y) v_\epsilon(y) dy \right) \varphi^{(r)}(x) dx \\ &= (-1)^r \int_{-\infty}^{\infty} v_\epsilon(y) \left( \int_{-\infty}^{\infty} f(x+y) \varphi^{(r)}(x) dx \right) dy \\ &= \int_{-\infty}^{\infty} v_\epsilon(y) \left( \int_{-\infty}^{\infty} f^{(r)}(x+y) \varphi(x) dx \right) dy \\ &= \int_{-\infty}^{\infty} \left( \int_{-\infty}^{\infty} f^{(r)}(x+y) v_\epsilon(y) dy \right) \varphi(x) dx \\ &= \left\langle \int_{-\infty}^{\infty} f^{(r)}(x+y) v_\epsilon(y) dy, \varphi(x) \right\rangle. \end{aligned}$$

So we have proved (3).

For all  $x \in \mathbb{R}$ , clearly,

$$|F_\epsilon^{(r)}(x)| \leq \|f^{(r)}(x + \cdot)\|_\Phi \|v_\epsilon\|_{(\bar{\Phi})} = \|f^{(r)}\|_\Phi.$$

Now we prove continuity of  $F_\epsilon^{(r)}(x)$  on  $\mathbb{R}$  ( $0 \leq r \leq n$ ). We show this for  $r = 0$  by contradiction: Assume that for some  $\epsilon > 0$ , point  $x^0$ , and subsequence  $|t_k| \rightarrow 0$

$$\left| \int_{-\infty}^{\infty} (f(x^0 + t_k + y) - f(x^0 + y)) v_\epsilon(y) dy \right| \geq \epsilon, \quad k \geq 1. \quad (4)$$

Since  $f \in L_\Phi$  we get easily  $f \in L_{1,loc}(\mathbb{R})$ . Then for any  $n = 1, 2, \dots$ ,  $f(t_k + y) \rightarrow f(y)$  in  $L_1(-n, n)$ . Therefore, there exists a subsequence, denoted again by  $\{t_k\}$ , such that  $f(t_k + y) \rightarrow f(y)$  a.e. in  $(-n, n)$ . Therefore, there exists a subsequence (for simplicity of notation we assume that it is coincident with  $\{t_k\}$ ) such that  $f(x^0 + t_k + y) \rightarrow f(x^0 + y)$  a.e. in  $(-\infty, \infty)$ .

On the other hand, without loss of generality we may assume that  $\rho(2f, \Phi) < \infty$ . Therefore by the Young inequality we get

$$\begin{aligned} & |f(x^0 + t_k + y) - f(x^0 + y)| |v_\epsilon(y)| \\ & \leq \Phi(|f(x^0 + t_k + y) - f(x^0 + y)|) + \bar{\Phi}(|v_\epsilon(y)|) \\ & \leq \frac{1}{2} \Phi(2|f(x^0 + y)|) + \frac{1}{2} \Phi(2|f(x^0 + t_k + y)|) + \bar{\Phi}(|v_\epsilon(y)|). \end{aligned}$$

The last expression belongs to  $L_1(\mathbb{R})$ , therefore by Lebesgue's theorem we have

$$\lim_{k \rightarrow \infty} \int_{-\infty}^{\infty} |f(x^0 + t_k + y) - f(x^0 + y)| |v_\epsilon(y)| dy = 0,$$

which contradicts (4). The cases  $1 \leq r \leq n$  are proved similarly. The continuity of  $F_\epsilon^{(r)}(x)$  has been proved.

The functions  $F_\epsilon^{(r)}(x)$  are continuous and bounded on  $\mathbb{R}$ . Therefore, it follows from the Kolmogorov inequality and (2)–(3) that

$$\begin{aligned} (\|f^{(k)}\|_\Phi - \epsilon)^n & \leq |F_\epsilon^{(k)}(0)|^n \leq \|F_\epsilon^{(k)}\|_\infty^n \\ & \leq C_{k,n} \|F_\epsilon\|_\infty^{n-k} \|F_\epsilon^{(n)}\|_\infty^k. \end{aligned} \quad (5)$$

On the other hand,

$$\|F_\epsilon\|_\infty \leq \|f(x + y)\|_\Phi \|v_\epsilon(y)\|_{(\bar{\Phi})} = \|f\|_\Phi, \quad (6)$$

$$\|F_\epsilon^{(n)}\|_\infty \leq \|f^{(n)}(x + y)\|_\Phi \|v_\epsilon(y)\|_{(\bar{\Phi})} = \|f^{(n)}\|_\Phi. \quad (7)$$

Combining (5)–(7), we get

$$(\|f^{(k)}\|_{\Phi} - \epsilon)^n \leq C_{k,n} \|f\|_{\Phi}^{n-k} \|f^{(n)}\|_{\Phi}^k.$$

By letting  $\epsilon \rightarrow 0$  we have (1).

To complete the proof, it remains to show that  $f^{(k)} \in L_{\Phi}(\mathbb{R})$ ,  $0 < k < n$  if  $f, f^{(n)} \in L_{\Phi}(\mathbb{R})$ .

Let  $\psi_{\lambda}(x) \in C_0^{\infty}(\mathbb{R})$ ,  $\psi_{\lambda}(x) \geq 0$ ,  $\psi_{\lambda}(x) = 0$  for  $|x| \geq \lambda$  and  $\int \psi_{\lambda}(x) dx = 1$ . We put  $f_{\lambda} = f * \psi_{\lambda}$ . Then  $f_{\lambda} \in C^{\infty}(\mathbb{R})$  because of  $f \in L_{1,loc}(\mathbb{R})$ . Therefore,  $f_{\lambda}^{(k)} = f * \psi_{\lambda}^{(k)}$ ,  $k \geq 0$ , and it is easy to check that  $f_{\lambda}^{(n)} = f^{(n)} * \psi_{\lambda}$ .

On the other hand, it follows from Lemma 2 that  $f_{\lambda}^{(k)} = f * \psi_{\lambda}^{(k)} \in L_{\Phi}(\mathbb{R})$ ,  $k \geq 0$ . Therefore, by the fact proved above, we have

$$\|f_{\lambda}^{(k)}\|_{\Phi}^n \leq C_{k,n} \|f_{\lambda}\|_{\Phi}^{n-k} \|f_{\lambda}^{(n)}\|_{\Phi}^k, \quad 0 < k < n.$$

Therefore, since

$$\|f_{\lambda}\|_{\Phi} \leq \|f\|_{\Phi} \cdot \|\psi_{\lambda}\|_1 = \|f\|_{\Phi}, \quad \|f_{\lambda}^{(n)}\|_{\Phi} \leq \|f^{(n)}\|_{\Phi} \|\psi_{\lambda}\|_1 = \|f^{(n)}\|_{\Phi}$$

we get that, for any  $0 \leq k \leq n$ , the sequence  $f_{\lambda}^{(k)}$  is bounded in  $L_{\Phi}(\mathbb{R})$ . Now we prove that, for any  $0 \leq k \leq n$ , there exists a subsequence, which is  $*$ -convergent to some  $g_k \in L_{\Phi}(\mathbb{R})$ . (We say that  $h_{\lambda}$  is  $*$ -convergent to  $h$ , where  $h_{\lambda}, h \in L_{\Phi}(\mathbb{R})$ , if  $\int h_{\lambda} v \rightarrow \int h v$  for all  $v \in L_{\Phi}(\mathbb{R})$ .) We will show, for example, the fact that  $f_{\lambda}$  is  $*$ -convergent to  $f$  by contradiction: Assume that for some  $\epsilon_0 > 0$ ,  $v \in L_{\Phi}(\mathbb{R})$  and a subsequence  $\lambda_k \rightarrow 0$ ,

$$\left| \int (f_{\lambda_k}(x) - f(x)) v(x) dx \right| \geq \epsilon_0, \quad k \geq 1. \quad (8)$$

Then, it is known that  $f_{\lambda} \rightarrow f$ ,  $\lambda \rightarrow 0$  in  $L_{1,loc}(\mathbb{R})$ . Therefore, there exists a subsequence  $\{k_m\}$  (for simplicity we assume that  $k_m = m$ ) such that  $f_{\lambda_k}(x) \rightarrow f(x)$  a.e.

We may assume that  $\rho(2f, \Phi) < \infty$ . Then it follows from Young's inequality that

$$|f_{\lambda}(x) - f(x)| |v(x)| \leq \frac{1}{2} \Phi(2|f_{\lambda}(x)|) + \frac{1}{2} \Phi(2|f(x)|) + \bar{\Phi}(|v(x)|),$$

moreover, the right side of the last inequality belongs to  $L_1(\mathbb{R})$ . Therefore, by virtue of Lebesgue's theorem we get

$$\lim_{k \rightarrow \infty} \int |f_{\lambda_k}(x) - f(x)| |v(x)| dx = 0$$

because of  $f_{\lambda_k}(x) \rightarrow f(x)$  a.e., which contradicts (8).

Finally, it follows from  $*$ -convergence  $f_\lambda \rightarrow f$  that for any  $\varphi \in C_0^\infty(\mathbb{R})$

$$\begin{aligned}\langle f_\lambda^{(k)}(x), \varphi(x) \rangle &= (-1)^k \langle f_\lambda(x), \varphi^{(k)}(x) \rangle \\ &\rightarrow (-1)^k \langle f(x), \varphi^{(k)}(x) \rangle = \langle f^{(k)}(x), \varphi(x) \rangle.\end{aligned}$$

Therefore, since the  $*$ -convergence of some subsequence of  $\{f_\lambda^{(k)}\}$  to  $g_k \in L_\Phi(\mathbb{R})$ , we get  $f^{(k)} = g_k \in L_\Phi(\mathbb{R})$  ( $0 < k < n$ ). So we have proved the fact that  $f^{(k)} \in L_\Phi(\mathbb{R})$  for all  $0 < k < n$  if  $f, f^{(n)} \in L_\Phi(\mathbb{R})$ . The proof is complete.

*Remark 1.* To obtain Theorem 1 we have developed the Stein method because, for example, the property  $[g(x+h) - g(x)]/h \rightarrow g'(x)$  in the  $L_p$  mean ( $1 \leq p < \infty$ ), which is used in [2], holds for  $L_\Phi$  only if  $\Phi(t)$  satisfies the  $\Delta_2$ -condition (see [10]).

*Remark 2.* For periodic functions we have:

**THEOREM 2.** Let  $\Phi(t)$  be an arbitrary Young function,  $f(x)$  and its generalized derivative  $f^{(n)}(x)$  be in  $L_\Phi(\mathbb{T})$ . Then  $f^{(k)}(x) \in L_\Phi(\mathbb{T})$  for all  $0 < k < n$  and

$$\| \| f^{(k)} \| \|_\Phi^n \leq C_{k,n} \| \| f \| \|_\Phi^{n-k} \| \| f^{(n)} \| \|_\Phi^k,$$

where  $\mathbb{T}$  is the torus and  $\| \|_\Phi$  is the corresponding norm.

*Remark 3.* By the representation [11, p. 135]

$$\| u \|_{(\Phi)} = \sup_{\| v \|_\Phi \leq 1} \left| \int u(x) v(x) dx \right|,$$

it is easy to see that the obtained results still hold for any Luxemburg norm.

## REFERENCES

1. A. N. Kolmogorov, On inequalities between upper bounds of the successive derivatives of an arbitrary function on an infinite interval, in "Amer. Math. Soc. Transl. Ser. 1," Vol. 2, pp. 233–243, Amer. Math. Soc., Providence, 1962.
2. E. M. Stein, Functions of exponential type, *Ann. Math.* **65** (1957), 582–592.
3. M. W. Certain and T. G. Kurtz, Landau-Kolmogorov inequalities for semigroups and groups, *Proc. Amer. Math. Soc.* **63** (1977), 226–230.
4. V. M. Tikhomirov and G. G. Magaril-Il'jaev, Inequalities for derivatives, in "Kolmogorov A. N. Selected Papers," pp. 387–390, Nauka, Moscow, 1985.
5. Ha Huy Bang, Some imbedding theorems for the spaces of infinite order of periodic functions, *Math. Notes* **43** (1988), 509–517.

6. Ha Huy Bang, On imbedding theorems for Sobolev spaces of infinite order, *Mat. Sb.* **178** (1988), 115–127.
7. Ha Huy Bang, Imbedding theorems for Sobolev spaces of infinite order, *Acta Math. Vietnam.* **14** (1989), 17–29.
8. Ha Huy Bang, A property of infinitely differentiable functions, *Proc. Amer. Math. Soc.* **108** (1990), 73–76.
9. R. O’Neil, Fractional integration in Orlicz space, I, *Trans. Amer. Math. Soc.* **115** (1965), 300–328.
10. W. Luxemburg, “Banach Function Spaces,” Thesis, Technische Hogeschool te Delft, The Netherlands, 1955.
11. M. A. Krasnoselsky and Y. B. Rutisky, “Convex Functions and Orlicz Spaces,” GITTL, Moscow, 1958; English translation, Noordhoff, Groningen, 1961.
12. R. Adams, “Sobolev Spaces,” Academic Press, New York, 1975.